

q-Partition Algebra Combinatorics

Tom Halverson

Department of Mathematics
Macalester College
Saint Paul, MN 55105
halverson@macalester.edu

Nathaniel Thiem

Department of Mathematics
University of Colorado at Boulder
Boulder, CO 80309
nathaniel.thiem@colorado.edu

September 8, 2009

Abstract

We study a q -analog $Q_r(n, q)$ of the partition algebra $P_r(n)$. The algebra $Q_r(n, q)$ arises as the centralizer algebra of the finite general linear group $\mathrm{GL}_n(\mathbb{F}_q)$ acting on a vector space $\mathcal{IR}_q^r$ coming from r -iterations of Harish-Chandra restriction and induction. For $n \geq 2r$, we show that $Q_r(n, q)$ has the same semisimple matrix structure as $P_r(n)$. We compute the dimension $d_{n,r}(q) = \dim(\mathcal{IR}_q^r)$ to be a q -polynomial that specializes as $d_{n,r}(1) = n^r$ and $d_{n,r}(0) = B(r)$, the r th Bell number. Our method is to write $d_{n,r}(q)$ as a sum over integer sequences which are q -weighted by inverse major index. We then find a basis of $\mathcal{IR}_q^r$ indexed by n -restricted q -set partitions of $\{1, \dots, r\}$ and show that there are $d_{n,r}(q)$ of these.

Introduction

The general linear group $\mathrm{GL}_n(\mathbb{C})$ and the symmetric group S_r both act on tensor space $V^{\otimes r}$, where V is the natural n dimensional representation of $\mathrm{GL}_n(\mathbb{C})$ and S_r acts by tensor place permutations. Classical Schur–Weyl duality says that these actions commute and that each action generates the full centralizer of the other, so that as a $(\mathrm{GL}_n(\mathbb{C}), S_r)$ -bimodule the tensor space has a multiplicity-free decomposition given by $V^{\otimes r} \cong \bigoplus_{\lambda} L(\lambda) \otimes S_r^{\lambda}$, where the $L(\lambda)$ are irreducible $\mathrm{GL}_n(\mathbb{C})$ -modules and the S_r^{λ} are the irreducible S_r -modules.

If we restrict $\mathrm{GL}_n(\mathbb{C})$ to its subgroup of orthogonal matrices $\mathrm{O}_n(\mathbb{C})$, then the centralizer algebra is Brauer’s centralizer algebra $B_r(n) = \mathrm{End}_{\mathrm{O}_n(\mathbb{C})}(V^{\otimes r})$. If we restrict further to the symmetric groups $S_{n-1} \subseteq S_n \subseteq \mathrm{O}_n(\mathbb{C}) \subseteq \mathrm{GL}_n(\mathbb{C})$, then the centralizer algebras are the partition algebras $P_r(n) = \mathrm{End}_{S_n}(V^{\otimes r})$ and $P_{r+\frac{1}{2}}(n) = \mathrm{End}_{S_{n-1}}(V^{\otimes r})$. Furthermore, the containments reverse:

$$\begin{array}{ccccccc} \text{subgroup } G \subseteq \mathrm{GL}_n(\mathbb{C}) : & S_{n-1} & \subseteq & S_n & \subseteq & \mathrm{O}_n(\mathbb{C}) & \subseteq & \mathrm{GL}_n(\mathbb{C}) \\ & \updownarrow & & \updownarrow & & \updownarrow & & \\ \text{centralizer algebra } \mathrm{End}_G(V^{\otimes r}) : & P_{r+\frac{1}{2}}(n) & \supseteq & P_r(n) & \supseteq & B_r(n) & \supseteq & \mathbb{C}S_r. \end{array}$$

The Brauer algebras were introduced in 1937 by Richard Brauer. The partition algebras arose early in the 1990s in the work of Martin [Mar1], [Mar2] and later, independently, in the work of Jones [Jo] (see also [HR]). For $r \in \frac{1}{2}\mathbb{Z}_{>0}$, the partition algebra $P_r(n)$ has a basis indexed by the set partitions of $\{1, 2, \dots, 2r\}$ and a multiplication given by “diagram multiplication.”

If the groups $\mathrm{GL}_n(\mathbb{C})$ and $\mathrm{O}_n(\mathbb{C})$ are replaced by their quantum groups $U_q(\mathfrak{gl}_n)$ and $U_q(\mathfrak{o}_n)$, respectively, then their centralizer algebras become

$$\begin{array}{llll} \text{quantum group :} & U_q(\mathfrak{o}_n) & \subseteq & U_q(\mathfrak{gl}_n) \\ & \updownarrow & & \updownarrow \\ \text{centralizer algebra :} & BMW_r(n, q) & \supseteq & H_r(q). \end{array}$$

The algebra $H_r(q)$ is the Iwahori–Hecke algebra of the symmetric group S_r , and its action on tensor space $V^{\otimes r}$ is due to Jimbo [Ji]. The algebra $BMW_r(n, q)$ is a q -analog of the Brauer algebra, called the Birman–Murakami–Wenzl algebra [BW], [Mur]. This leads naturally to the problem of finding a q -analog of the partition algebra. One might consider replacing the symmetric group S_n with its Iwahori–Hecke algebra $H_n(q)$, but $H_n(q)$ does not have a Hopf coproduct allowing it to act on the tensor product representation $V^{\otimes r}$. Furthermore, the Hecke algebra $H_n(q)$ is not found as a subalgebra of $U_q(\mathfrak{o}_n)$ (in fact, even the containment of $U_q(\mathfrak{o}_n)$ is $U_q(\mathfrak{gl}_n)$ is more subtle than $\mathrm{O}_n(\mathbb{C}) \subseteq \mathrm{GL}_n(\mathbb{C})$).

In this paper, we take a different approach to defining a q -partition algebra in which we replace the underlying tensor space with a module constructed by iterations of restriction and induction of finite general linear group modules. This approach was first proposed in unpublished work of T. Halverson and A. Ram and is motivated by the analogous construction of the partition algebra through restriction and induction of symmetric group modules (see [HR]). A forthcoming paper by T. Halverson, A. Ram, and N. Thiern will further study the q -partition algebra, and the analysis of the underlying restriction-induction module found in this paper is essential to that work.

The paper is organized as follows:

1. In Section 1, we describe the construction of a q -partition algebra $Q_r(n, q)$ as the centralizer of the general linear $\mathrm{GL}_n(\mathbb{F}_q)$ over a finite field $\mathbb{F}_q$ having q elements. For $n \geq 2r$, $Q_r(n, q)$ is the centralizer of $\mathrm{GL}_n(\mathbb{F}_q)$ acting a vector space $\mathcal{IR}_q^r$, consisting of r iterations of Harish-Chandra restriction and induction. At $q = 1$, we have $\mathcal{IR}_1^r \cong V^{\otimes r}$, and we think of the symmetric group S_n as the $q \rightarrow 1$ limit of $\mathrm{GL}_n(\mathbb{F}_q)$. We show that $Q_r(n, q)$ and $P_r(n, q)$ each have dimension equal to the Bell number $B(2r)$, and that for $n \geq 2r$ they have the same matrix block structure as semisimple algebras.
2. In Section 2, we combinatorially study the dimension of $\mathcal{IR}_q^k$. We show that

$$\dim(\mathcal{IR}_q^r) = d_{n,r}(q) = \sum_{\ell=1}^n S(r, \ell) [n] [n-1] \cdots [n-\ell+1], \quad (0.1)$$

where $S(r, \ell)$ is a Stirling number of the second kind and $[j] = (q^j - 1)/(q - 1)$ is a q -integer. The q -polynomial $d_{n,r}(q)$ that appears in this formula has the property that $d_{n,r}(1) = n^r$ and $d_{n,r}(0) = B(r)$, the r th Bell number or number of partitions of $\{1, \dots, r\}$ into subsets. Thus $d_{n,r}(q)$ is a q -analog of both n^r and $B(r)$, and it interpolates between the two as q ranges through $0 \leq q \leq 1$. Our method is to write the dimension as a q -weighted sum over sequences $a = (a_1, \dots, a_r)$, where each sequence is weighted by an analog of the inverse major index $\mathrm{imaj}(a)$. This is done in Proposition 2.3. We then use a Schensted bijection (see (2.1)) and the decomposition of $\mathcal{IR}_q^r$ as a $(\mathrm{GL}_n(\mathbb{F}_q), Q_r(n, q))$ -bimodule to prove formula (0.1).

3. In Section 3, we define (n -restricted) q -set partitions of $\{1, \dots, r\}$, and we show that $d_{n,r}(q)$ enumerates these objects. We study the module $\mathcal{IR}_q^r$ and find a basis for it that is indexed by these q -weighted set partitions of $\{1, \dots, r\}$.

T. Halverson was partially supported by the National Science Foundation under grant DMS-0100975. This research was completed while the authors were in residence at the Mathematical Sciences Research Institute (MSRI) in Spring 2008 for the program in Combinatorial Representation Theory. We are grateful for the support and the stimulating research environment at MSRI. We thank Arun Ram for numerous useful conversations, and we thank Vic Reiner and Dennis Stanton for a helpful conversation about the distribution of the statistics inv and maj used in the proof of Proposition 2.3. We are grateful to the anonymous referees for helpful suggestions.

1 A q -Partition Algebra

View $S_{n-1} \subseteq S_n$ under the natural embedding. The n -dimensional permutation module V for S_n is isomorphic to the induced module, $V \cong \text{Ind}_{S_{n-1}}^{S_n}(\mathbb{1}_{n-1}) = \mathbb{C}S_n \otimes_{S_{n-1}} \mathbb{1}_{n-1}$, where $\mathbb{1}_{n-1}$ is the trivial S_{n-1} -module. In [HR], Halverson and Ram emphasize viewing tensor products of S_n -modules via restriction and induction and the “tensor identity,”

$$\begin{aligned} \text{Ind}_{S_{n-1}}^{S_n} \text{Res}_{S_{n-1}}^{S_n}(W) &\cong W \otimes V \\ g \otimes_{S_{n-1}} w &\mapsto gm \otimes (g \otimes_{S_{n-1}} \mathbb{1}_{n-1}), \end{aligned} \quad (1.1)$$

where W is any S_n module, $w \in W, g \in S_{n-1}$. For $r \in \mathbb{Z}_{\geq 0}$ define,

$$\mathcal{IR}_1^r = \underbrace{\text{Ind}_{S_{n-1}}^{S_n} \text{Res}_{S_{n-1}}^{S_n} \cdots \text{Ind}_{S_{n-1}}^{S_n} \text{Res}_{S_{n-1}}^{S_n}}_{2r \text{ functors}}(\mathbb{1}_n), \quad (1.2)$$

and

$$\mathcal{IR}_1^{r+\frac{1}{2}} = \underbrace{\text{Res}_{S_{n-1}}^{S_n} \text{Ind}_{S_{n-1}}^{S_n} \text{Res}_{S_{n-1}}^{S_n} \cdots \text{Ind}_{S_{n-1}}^{S_n} \text{Res}_{S_{n-1}}^{S_n}}_{2r+1 \text{ functors}}(\mathbb{1}_n), \quad (1.3)$$

Then, by induction, $V^{\otimes r} \cong \mathcal{IR}_1^r$, and therefore the partition algebra satisfies, for $n \geq 2r$,

$$\begin{aligned} P_r(n) &= \text{End}_{S_n}(V^{\otimes r}) \cong \text{End}_{S_n}(\mathcal{IR}_1^r), \\ P_{r+\frac{1}{2}}(n) &= \text{End}_{S_{n-1}}(V^{\otimes r}) \cong \text{End}_{S_{n-1}}(\mathcal{IR}_1^{r+\frac{1}{2}}). \end{aligned} \quad (1.4)$$

Key to the decomposition of $\mathcal{IR}_1^r$ into irreducible symmetric group modules are the restriction and induction rules

$$\text{Res}_{S_{n-1}}^{S_n}(S_n^\lambda) = \bigoplus_{\mu=\lambda-\square} S_{n-1}^\mu \quad \text{and} \quad \text{Ind}_{S_{n-1}}^{S_n}(S_{n-1}^\mu) = \bigoplus_{\lambda=\mu+\square} S_n^\lambda, \quad (1.5)$$

where S_n^λ is the irreducible S_n module labeled by the partition $\lambda \vdash n$, S_{n-1}^μ is the irreducible S_{n-1} module labeled by the partition $\mu \vdash (n-1)$, and $\lambda - \square$ and $\mu + \square$ denote adding and removing a box from the partition, respectively.

One can view the symmetric group S_n as the $q \rightarrow 1$ limit of the general linear group $G_n = \text{GL}_n(\mathbb{F}_q)$ over the finite field $\mathbb{F}_q$. Indeed, if B is the Borel subgroup of upper triangular matrices in $\text{GL}_n(\mathbb{F}_q)$, then

$$|\text{GL}_n(\mathbb{F}_q)/B| = [n][n-1] \cdots [2][1],$$

where $[n] = 1 + q + \dots q^n$ is a q -analog of n so that $[n][n-1]\dots[2][1]$ is a q -analog of $n!$. Furthermore, the irreducible unipotent representations of G_n are denoted G_n^λ and labeled by partitions $\lambda \vdash n$ (see, for example, [Mac, §4.3]).

We view $G_{n-1} \subseteq G_n$ as a Levi subgroup with blocks of size 1 and $n-1$ (see Section 3.3). Under Harish-Chandra restriction Resf and induction Indf (see Section 3.3) these modules satisfy exactly the same rules as (1.5), namely (see [Mac, §4.3]),

$$\text{Resf}_{G_{n-1}}^{G_n}(G_n^\lambda) = \bigoplus_{\mu=\lambda-\square} G_{n-1}^\mu \quad \text{and} \quad \text{Indf}_{G_{n-1}}^{G_n}(G_{n-1}^\mu) = \bigoplus_{\lambda=\mu+\square} G_n^\lambda. \quad (1.6)$$

For $r \in \mathbb{Z}_{\geq 0}$ define,

$$\mathcal{IR}_q^r = \underbrace{\text{Indf}_{G_{n-1}}^{G_n} \text{Resf}_{G_{n-1}}^{G_n} \cdots \text{Indf}_{G_{n-1}}^{G_n} \text{Resf}_{G_{n-1}}^{G_n}}_{2r \text{ functors}}(\mathbb{1}_{G_n}), \quad (1.7)$$

and

$$\mathcal{IR}_q^{r+\frac{1}{2}} = \underbrace{\text{Res}_{G_{n-1}}^{G_n} \text{Ind}_{G_{n-1}}^{G_n} \text{Res}_{G_{n-1}}^{G_n} \cdots \text{Ind}_{G_{n-1}}^{G_n} \text{Res}_{G_{n-1}}^{G_n}}_{2r+1 \text{ functors}}(\mathbb{1}_{G_n}), \quad (1.8)$$

Then, for integers $n \geq 2r$, define

$$\begin{aligned} Q_r(n, q) &= \text{End}_{G_n}(\mathcal{IR}_q^r), \\ Q_{r+\frac{1}{2}}(n, q) &= \text{End}_{G_{n-1}}(\mathcal{IR}_q^{r+\frac{1}{2}}). \end{aligned} \quad (1.9)$$

Equation (1.9) completely defines $Q_r(n, q)$ as an algebra of endomorphisms, however considerable work needs to be done to find a natural set of generators for $Q_r(n, q)$ and the relations that they satisfy. This will be the subject of a forthcoming paper by T. Halverson, A. Ram, and N. Thiern. The analysis in this paper will be foundational to that work.

The *Bratteli diagram* $\mathfrak{B}_n$ is a graph that encodes the decomposition of $\mathcal{IR}_q^r$. Let $\mathfrak{B}_n$ have vertices organized into levels indexed by $r \in \frac{1}{2}\mathbb{Z}_{\geq 0}$ such that the vertices on level r are labeled by the set of integer partitions Λ_n^r defined by

$$\begin{aligned} \Lambda_n^r &= \{ \lambda = (\lambda_1, \dots, \lambda_t) \vdash n \mid \lambda_2 + \dots + \lambda_t \leq r \}, \\ \Lambda_n^{r+\frac{1}{2}} &= \{ \mu = (\mu_1, \dots, \mu_t) \vdash (n-1) \mid \mu_2 + \dots + \mu_t \leq r \}. \end{aligned} \quad (1.10)$$

There is an edge in $\mathfrak{B}_n$ from $\lambda \in \Lambda_n^r$ to $\mu \in \Lambda_n^{r+\frac{1}{2}}$ or $\mu \in \Lambda_n^{r-\frac{1}{2}}$ if and only if $\lambda = \mu + \square$. For example, the Bratteli diagram $\mathfrak{B}_6$, for $0 \leq r \leq 3$, is shown in Figure 1. The edges in $\mathfrak{B}_n$ describe the restriction and induction rules in (1.5) and (1.6). Since Λ_n^0 contains only the partition (n) , which labels the trivial G_n or S_n -module, the Bratteli diagram has the property that the vertices on level r label the irreducible G_n modules which appear in $\mathcal{IR}_q^r$ or, equivalently, the irreducible S_n modules which appear in $\mathcal{IR}^r$. Furthermore, the number of paths from the top of the diagram to $\lambda \in \Lambda_n^r$ is the multiplicity of G_n^λ in $\mathcal{IR}_q^r$. The number of these paths is also indicated below each vertex in Figure 1.

From double-centralizer theory (see for example [HR, Theorem 5.4]) we have the following properties which justify calling $Q_r(n, q)$ a q -analog of the partition algebra $P_r(n)$. These results follow from the fact that the decomposition of $\mathcal{IR}_q^r$ as a G_n module is the same as the decomposition of $\mathcal{IR}_1^r$ as an S_n -module. We assume $r, n \in \mathbb{Z}_{\geq 0}$ and $n \geq 2r$,




Figure 1: The Bratteli diagram $\mathfrak{B}_6$ for $0 \leq r \leq 3$. The rows contain the partitions in Λ_6^r , the paths from the top of the diagram to $\lambda \in \Lambda_6^r$ are the r -vacillating tableaux of shape λ , and the number of paths to λ is m_r^λ . The label on vertex λ is m_r^λ .

1. The irreducible representations of $P_r(n)$ and $Q_r(n, q)$ are each indexed by the partitions in Λ_n^r . These are the partitions which label the irreducible S_n and G_n modules which appear in $\mathcal{IR}_1^r$ and $\mathcal{IR}_q^r$, respectively.
2. As bimodules for $(S_n, P_r(n))$ and $(G_n, Q_r(n, q))$, we have

$$\mathcal{IR}_1^r \cong \bigoplus_{\lambda \in \Lambda_n^r} S_n^\lambda \otimes P_r^\lambda \quad \text{and} \quad \mathcal{IR}_q^r \cong \bigoplus_{\lambda \in \Lambda_n^r} G_n^\lambda \otimes Q_r^\lambda, \quad (1.11)$$

where P_r^λ and Q_r^λ are irreducible $P_r(n)$ and $Q_r(n, q)$ -modules, respectively.

3. $m_r^\lambda = \dim(P_r^\lambda) = \dim(Q_r^\lambda)$ equals the multiplicity of S_n^λ in $\mathcal{IR}^r$ (or, equivalently, the multiplicity of G_n^λ in $\mathcal{IR}_q^r$). The value of m_r^λ can be computed by counting paths to λ in the Bratteli diagram. Each of these paths is a sequence of partitions obtained by iteratively removing and adding a box; these are called r -vacillating tableaux of shape λ (see Section 2) and they are also studied in [CDDSY] and [HL].
4. By general Wedderburn theory, $\dim(P_r(n)) = \sum_{\lambda \in \Lambda_n^r} \dim(P_r^\lambda)^2 = \sum_{\lambda \in \Lambda_n^r} \dim(Q_r^\lambda)^2 = \dim(Q_r(n, q))$. It follows that $\dim(P_r(n)) = \dim(Q_r(n, q))$ equals the number of set partitions of $\{1, 2, \dots, 2r\}$ which is the Bell number $B(2r)$.
5. For $r \in \mathbb{Z}_{>0}$, there is a natural chain of embeddings

$$Q_{r-\frac{1}{2}}(n, q) \subseteq Q_r(n, q) \subseteq Q_{r+\frac{1}{2}}(n, q),$$

The restriction rule for $Q_{r-\frac{1}{2}}(n, q) \subseteq Q_r(n, q)$ is given by the Bratteli diagram. Namely, the module $Q_{r-\frac{1}{2}}^\mu$ is a component of $\text{Res}_{Q_{r-\frac{1}{2}}(n, q)}^{Q_r(n, q)}(Q_r^\lambda)$ if and only if μ and λ are connected by an edge in $\mathfrak{B}_n$. This same rule holds if $Q_r(n, q)$ is replaced by $P_r(n)$.

Since restriction from S_n to S_{n-1} preserves the dimension of an S_n -module, and induction from S_{n-1} to S_n increases the dimension by a factor of n , we have that $\dim(\mathcal{IR}_1^r) = n^r$. This is to be expected, since $\mathcal{IR}_1^r \cong V^{\otimes r}$ and $\dim(V) = n$. The dimension of the irreducible symmetric group modules are $\dim(S_n^\lambda) = f_n^\lambda$, the number of standard Young tableaux of shape λ (given by the hook formula). The dimensions of the irreducible partition algebra modules are $\dim(P_r^\lambda) = m_r^\lambda$, the number of r -vacillating tableau of shape λ . Computing dimensions on both sides of the first equation in (1.11) gives the identity

$$n^r = \sum_{\lambda \vdash n} f_n^\lambda m_r^\lambda. \quad (1.12)$$

A combinatorial proof of (1.12) is given in [HL] by defining a Schensted-like insertion/deletion process to find a bijection

$$\{(a_1, \dots, a_r)\} \longleftrightarrow \{(P, Q)\}$$

between integer sequences $(a_1, \dots, a_r)$ with $a_j \in \{1, \dots, n\}$ and pairs (P, Q) , where P is a standard Young tableau of shape $\lambda \vdash n$ and Q is an r -vacillating tableau of shape $\lambda \vdash n$. If we compute the dimensions on each side of the second equation in (1.11) we get

$$d_{n,r}(q) = \dim(\mathcal{IR}_q^r) = \sum_{\lambda} f_n^\lambda(q) m_k^\lambda, \quad (1.13)$$

where now $f_n^\lambda(q) = \dim(G_n^\lambda)$ is the well-known q -analog of f_n^λ , given by the q -hook formula and $d_{n,r}(q)$ is a q -polynomial which satisfies $d_{n,r}(1) = n^r$. In Section 2, we compute this dimension explicitly by carrying the q -weights from $f^\lambda(q)$ across the Schensted insertion.

2 Combinatorial Computation of $d_{n,r}(q)$

This section gives a purely combinatorial derivation of the formula for the q -polynomial $d_{n,r}(q)$. In Section 3, we give a basis of $\mathcal{IR}_q^r$ and we count the elements of the basis to give another proof that $\dim(\mathcal{IR}_q^r) = d_{n,r}(q)$.

2.1 The Delete-Insert Schensted Algorithm

For $n, r \in \mathbb{Z}_{>0}$, define

$$\{1, \dots, n\}^r = \{ (a_1, \dots, a_r) \mid a_j \in \{1, \dots, n\} \}.$$

This set has cardinality n^r . For a partition $\lambda \vdash n$, a standard tableau of shape λ is a filling of the boxes of the Young diagram of λ with integers $1, \dots, n$ such that the rows increase left-to-right and the columns increase top-to-bottom. As in [HL] we define an algorithm that maps sequences in $\{1, \dots, n\}^r$ to standard tableaux. Let $a = (a_1, a_2, \dots, a_r)$ and recursively define P_i and $P_{i+\frac{1}{2}}$ for $0 \leq i \leq r$, by

$$\begin{aligned} P_0 &= \boxed{1 \mid 2 \mid \cdots \mid n}, \\ P_{i+\frac{1}{2}} &= P_{i-1} \xrightarrow{\text{jdt}} a_i, \quad 0 \leq i \leq r-1, \\ P_{i+1} &= P_{i+\frac{1}{2}} \xleftarrow{\text{RSK}} a_i, \quad 0 \leq i \leq r-1, \end{aligned} \tag{2.1}$$

where this notation means that we first remove the letter a_i from P_i using Schützenberger's *jeu-de-taquin* to get a tableau $P_{i+\frac{1}{2}}$, and then we reinsert a_i into $P_{i+\frac{1}{2}}$ using *Robinson-Schensted-Knuth row insertion* to obtain P_{i+1} . See [Sta2, A1.2, 7.11] for the definitions of jeu-de-taquin and RSK insertion. Example 1 provides an example of the application of this algorithm.

For $0 \leq i \leq k$, let $\lambda^{(i)}$ be the partition shape of the tableau P_i and let $\lambda^{(i+\frac{1}{2})}$ be the partition shape of $P_{i+\frac{1}{2}}$. The final tableau $P_a = P_r$ that results from the insertion of $a = (a_1, \dots, a_r)$ is the *insertion tableau*. It is a standard Young tableaux of shape $\lambda = \lambda^{(r)}$. The sequence of shapes that arise along the way,

$$Q_a = \left((n) = \lambda^{(0)}, \lambda^{(\frac{1}{2})}, \lambda^{(1)}, \dots, \lambda^{(r)} = \lambda \right),$$

is the *recording tableau* of the sequence. The recording tableaux that appear in this process are uniquely described by the following properties:

1. $\lambda^{(0)} = (n)$,
2. For $0 \leq i \leq r-1$, $\lambda^{(i+\frac{1}{2})}$ is a partition of $n-1$ that is obtained from $\lambda^{(i)}$ by deleting a box,
3. For $1 \leq i \leq r$, $\lambda^{(i)}$ is a partition of n that is obtained from $\lambda^{(i-\frac{1}{2})}$ by adding a box.

If $\lambda = \lambda^{(r)}$, then a tableau satisfying these properties is called a *r-vacillating tableau of shape λ* . See [HL] and [CDDSY]. The partition shapes that appear in the ℓ th step in the process of inserting $a \in \{1, \dots, n\}^r$ are in the set

$$\begin{aligned} \Lambda_n^\ell &= \{ \lambda = (\lambda_1, \dots, \lambda_t) \vdash n \mid \lambda_2 + \cdots + \lambda_t \leq \ell \}, \\ \Lambda_n^{\ell+\frac{1}{2}} &= \{ \lambda = (\lambda_1, \dots, \lambda_t) \vdash n-1 \mid \lambda_2 + \cdots + \lambda_t \leq \ell \}. \end{aligned}$$

The r -vacillating tableau also appear in the Bratteli diagram $\mathfrak{B}_n$ shown in Figure 1 for $n = 6$ and $0 \leq r \leq 3$. The paths from the top of the diagram to λ on level r are the r -vacillating tableaux of shape λ , and m_r^λ is the number of r -vacillating tableaux of shape λ . When $r \geq 2n$, the number is independent of n . We refer to these paths as “tableaux” since they determine paths in the Bratteli diagram in the same way that standard Young tableaux determine paths in Young’s lattice.

We let $a \xrightarrow{\text{DI}} (P_a, Q_a)$ denote the “delete-insert” process defined in (2.1), which associates each $a \in \{1, \dots, n\}^r$ with a pair (P_a, Q_a) consisting of a standard tableau P_a and an r -vacillating tableau Q_a , each of shape $\lambda \in \Lambda_n^r$. In [HL] this algorithm is shown to be invertible and thus provides a bijection

$$\{1, \dots, n\}^r \xrightarrow{\text{DI}} \bigsqcup_{\lambda \in \Lambda_n^r} \left\{ (P, Q) \mid \begin{array}{l} P \text{ is a standard tableau of shape } \lambda \\ Q \text{ is a } r\text{-vacillating tableau of shape } \lambda \end{array} \right\}. \quad (2.2)$$

This gives a combinatorial proof of the identity

$$n^r = \sum_{\lambda \in \Lambda_n^r} f_n^\lambda m_r^\lambda, \quad (2.3)$$

where f_n^λ is the number of standard tableaux of shape λ (given by the hook formula), and m_r^λ is the number of r -vacillating tableaux of shape λ .

2.2 Delete/Insertion and Major Index

We now show that the bijection (2.1) carries the backsteps associated to integer sequences to the descent set on standard tableaux. We first map sequences in $\{1, \dots, n\}^r$ to permutations in S_n using following surjection

$$\begin{aligned} \{1, \dots, n\}^r &\rightarrow S_n \\ a = (a_1, \dots, a_r) &\mapsto w_a = \text{RT}(1, 2, \dots, n, a_1, \dots, a_r) \end{aligned} \quad (2.4)$$

where $\text{RT}(1, 2, \dots, n, a_1, \dots, a_r)$ is the permutation consisting of the rightmost occurrence of each integer in $\{1, \dots, n\}$. For example,

$$a = (2, 1, 3, 1, 6, 4, 6, 3, 4) \mapsto w_a = \text{RT}(1, 2, 3, 4, 5, 6, 2, 1, 3, 1, 6, 4, 6, 3, 4) = (5, 2, 1, 6, 3, 4).$$

Alternatively, we can produce $w_a = (b_1, b_2, \dots, b_n)$ iteratively using the following algorithm.

$$\begin{aligned} (1) \quad & w^{(0)} = (1, 2, \dots, n), \\ (2) \quad & w^{(i+1)} \text{ is obtained from } w^{(i)} \text{ by deleting } a_i \text{ from } w^{(i)} \\ & \text{and then appending } a_i \text{ to the right of } w^{(i)}, \\ (3) \quad & w_a = w^{(r)}. \end{aligned} \quad 1 \leq i < r. \quad (2.5)$$

Applying this algorithm to $a = (2, 1, 3, 1, 6, 4, 6, 3, 4)$, for example, yields the same w_a as above:

$$\begin{array}{ll} w^{(0)} = (1, 2, 3, 4, 5, 6) & w^{(5)} = (4, 5, 2, 3, 1, 6) \\ w^{(1)} = (1, 3, 4, 5, 6, 2) & w^{(6)} = (5, 2, 3, 1, 6, 4) \\ w^{(2)} = (3, 4, 5, 6, 2, 1) & w^{(7)} = (5, 2, 1, 6, 4, 3) \\ w^{(3)} = (4, 5, 6, 2, 1, 3) & w^{(8)} = (5, 2, 1, 6, 3, 4) \\ w^{(4)} = (4, 5, 6, 2, 3, 1) & w_a = (5, 2, 1, 6, 3, 4). \end{array}$$

It is clear that the processes defined in (2.4) and (2.5) yield the same result since the elements of a are cycled to the right end of w_a in the order that they appear in a .

The *backsteps* (see for example [Lo]) in a permutation $w = (w_1, w_2, \dots, w_n) \in S_n$ are

$$BS(w) = \{ i \mid i+1 \text{ is to the left of } i \text{ in } w = (w_1, w_2, \dots, w_n) \}. \quad (2.6)$$

The *descent set* in $w \in S_n$ is defined by $Des(w) = Des(w_1, w_2, \dots, w_n) = \{ i \mid w_i > w_{i+1} \}$, and it is easy to check that $BS(w) = Des(w^{-1})$. For example if $w = (5, 2, 1, 6, 3, 4)$ then $BS(w) = Des(w^{-1}) = \{1, 4\}$. If P is a standard tableau, then the descent set of P is

$$Des(P) = \{ i \mid i+1 \text{ is in a lower row than } i \text{ in } P \}. \quad (2.7)$$

For example $Des \left(\begin{array}{|c|c|c|c|} \hline 1 & 2 & 5 & 6 \\ \hline 3 & 7 & 9 & 10 \\ \hline 4 & 8 & & \\ \hline \end{array} \right) = \{2, 3, 6, 7\}$. See Example 1 for an illustration of the following proposition.

Proposition 2.1. *If $a \in \{1, \dots, n\}^r$ and $a \xrightarrow{\text{DI}} (P_a, Q_a)$, where P_a is a standard tableau of shape $\lambda \in \Lambda_n^r$ and Q_a is an r -vacillating tableau, then*

$$BS(w_a) = Des(P_a).$$

Proof. The proof is by induction on the length r of $a = (a_1, \dots, a_r)$. If $r = 0$, then $w = \emptyset$ and $P_a = \boxed{1} \boxed{2} \dots \boxed{n}$. In this case, $w_a = (1, 2, \dots, n)$ has $BS(w_a) = \emptyset = Des(P_a)$.

Now let $r > 0$ and $(a_1, \dots, a_{r-1}) \xrightarrow{\text{DI}} (P_{r-1}, Q_{r-1})$. Then $P_a = (P_{r-1} \xrightarrow{\text{jdt}} a_r) \xleftarrow{\text{RSK}} a_r$, and by induction $Des(P_{r-1}) = BS(w_{(a_1, \dots, a_{r-1})})$. By (2.5), the permutation $w_{(a_1, \dots, a_r)}$ is the same as $w_{(a_1, \dots, a_{r-1})}$ except that it has a_r moved to the rightmost position. Since a_r is now to the right of both $a_r - 1$ and $a_r + 1$, and this is the only change made, we know that

- (a) $(a_r - 1, a_r)$ is not a backstep in w_a ,
- (b) $(a_r, a_r + 1)$ is a backstep in w_a , and
- (c) all other $(i, i + 1)$ relationships are the same in w_a as they were in $w_{(a_1, \dots, a_{r-1})}$.

These same relationships happen in P :

- (a') When a_r is deleted from P_{r-1} (via jeu-de-taquin) and then reinserted (via RSK), it ends up in the first row of P . Thus $(a_r - 1, a_r)$ is not a descent in P .
- (b') If $a_r + 1$ was in the first row of P_{r-1} then a_r bumps it to a lower row. Otherwise, it was already in a lower row, and either way $(a_r, a_r + 1)$ is a descent in P .
- (c') Whenever i gets bumped into the next row, if $i + 1$ is in that row, i will bump $i + 1$ into a lower row. So if $(i, i + 1)$ is a descent it will remain a descent. If $(i, i + 1)$ is not a descent, then we must consider the case when $i + 1$ gets bumped lower than i . This only happens if i and $i + 1$ are in the same row. But in this case a number that might bump $i + 1$ would have to be lower than $i + 1$ and thus lower than i . So it might potentially bump i but it would not bump $i + 1$.

It follows by induction that $Des(P_a) = BS(w_a)$, as desired. $\square$

The inverse major index $imaj$ of a permutation $w \in S_n$ is the sum of the backsteps in w , and the major index maj of a standard tableau P is the sum of the descents in P . That is,

$$imaj(w) = \sum_{i \in BS(w)} i \quad \text{and} \quad maj(P) = \sum_{i \in Des(P)} i. \quad (2.8)$$

Note that the major index of w is $maj(w) = \sum_{i \in Des(w)} i$, and $imaj(w) = maj(w^{-1})$. Let q be an indeterminate (in Section 3 we will specialize q to be a prime power). For $\lambda \vdash n$, a q -analog of the hook number f_n^λ is given by

$$f_n^\lambda(q) = \sum_T q^{maj(T)}, \quad (2.9)$$

where the sum is over all standard tableaux T of shape λ . Then $f_n^\lambda(q)$ is the dimension of the irreducible unipotent $GL_n(\mathbb{F}_q)$ -module labeled by λ and it is also given by the q -hook formula (see [Mac, IV.6.7]).

Corollary 2.2. *For all $n, r \in \mathbb{Z}_{>0}$, we have*

$$\sum_{a \in \{1, \dots, n\}^r} q^{imaj(w_a)} = \sum_{\lambda \in \Lambda_n^r} \sum_{(P, Q)} q^{maj(P)} = \sum_{\lambda \in \Lambda_n^r} f_n^\lambda(q) m_r^\lambda,$$

where (P, Q) ranges over all pairs consisting of a standard tableau P of shape λ and an r -vacillating tableau Q of shape λ , and w_a is defined in (2.4).

Proof. The first equality follows immediately from the fact that the delete-insert bijection (2.2) pairs $a \in \{1, \dots, n\}^r$ with $\{(P_a, Q_a)\}$ and carries $q^{imaj(w_a)}$ to $q^{maj(P)}$. The second equality follows from (2.9) and from the fact that m_r^λ equals the number of r -vacillating tableaux Q of shape λ . $\square$

Example 1. The following table illustrates the process of delete-inserting the sequence $a = (3, 5, 2, 3, 2) \in \{1, \dots, 6\}^5$ to produce a pair (P_a, Q_a) of shape $\lambda = (2, 2, 1, 1)$. The reader should

observe that at each step in this process the backsteps in w_a equal the descents in P_a .

i	a_i	P_a	a	w_a	$BS(w_a) = Des(P_a)$										
0		<table><tr><td>1</td><td>2</td><td>3</td><td>4</td><td>5</td><td>6</td></tr></table>	1	2	3	4	5	6	$\emptyset$	$(1, 2, 3, 4, 5, 6)$	$\emptyset$				
1	2	3	4	5	6										
$\frac{1}{2}$	3	<table><tr><td>1</td><td>2</td><td>4</td><td>5</td><td>6</td></tr></table> $\xrightarrow{\text{jdt}} 3$	1	2	4	5	6								
1	2	4	5	6											
1		<table><tr><td>1</td><td>2</td><td>3</td><td>5</td><td>6</td></tr><tr><td>4</td><td></td><td></td><td></td><td></td></tr></table> $\xleftarrow{\text{RSK}} 3$	1	2	3	5	6	4					(3)	$(1, 2, 4, 5, 6, 3)$	$\{3\}$
1	2	3	5	6											
4															
$1\frac{1}{2}$	5	<table><tr><td>1</td><td>2</td><td>3</td><td>6</td></tr><tr><td>4</td><td></td><td></td><td></td></tr></table> $\xrightarrow{\text{jdt}} 5$	1	2	3	6	4								
1	2	3	6												
4															
2		<table><tr><td>1</td><td>2</td><td>3</td><td>5</td></tr><tr><td>4</td><td>6</td><td></td><td></td></tr></table> $\xleftarrow{\text{RSK}} 5$	1	2	3	5	4	6			$(3, 5)$	$(1, 2, 4, 6, 3, 5)$	$\{3, 5\}$		
1	2	3	5												
4	6														
$2\frac{1}{2}$	2	<table><tr><td>1</td><td>3</td><td>5</td></tr><tr><td>4</td><td>6</td><td></td></tr></table> $\xrightarrow{\text{jdt}} 2$	1	3	5	4	6								
1	3	5													
4	6														
3		<table><tr><td>1</td><td>2</td><td>5</td></tr><tr><td>3</td><td>6</td><td></td></tr><tr><td>4</td><td></td><td></td></tr></table> $\xleftarrow{\text{RSK}} 2$	1	2	5	3	6		4			$(3, 5, 2)$	$(1, 4, 6, 3, 5, 2)$	$\{2, 3, 5\}$	
1	2	5													
3	6														
4															
$3\frac{1}{2}$	3	<table><tr><td>1</td><td>2</td><td>5</td></tr><tr><td>4</td><td>6</td><td></td></tr></table> $\xrightarrow{\text{jdt}} 3$	1	2	5	4	6								
1	2	5													
4	6														
4		<table><tr><td>1</td><td>2</td><td>3</td></tr><tr><td>4</td><td>5</td><td></td></tr><tr><td>6</td><td></td><td></td></tr></table> $\xleftarrow{\text{RSK}} 3$	1	2	3	4	5		6			$(3, 5, 2, 3)$	$(1, 4, 6, 5, 2, 3)$	$\{3, 5\}$	
1	2	3													
4	5														
6															
$4\frac{1}{2}$	2	<table><tr><td>1</td><td>3</td></tr><tr><td>4</td><td>5</td></tr><tr><td>6</td><td></td></tr></table> $\xrightarrow{\text{jdt}} 2$	1	3	4	5	6								
1	3														
4	5														
6															
5		<table><tr><td>1</td><td>2</td></tr><tr><td>3</td><td>5</td></tr><tr><td>4</td><td></td></tr><tr><td>6</td><td></td></tr></table> $\xleftarrow{\text{RSK}} 2$	1	2	3	5	4		6		$(3, 5, 2, 3, 2)$	$(1, 4, 6, 5, 3, 2)$	$\{2, 3, 5\}$		
1	2														
3	5														
4															
6															

2.3 Set Partitions and Major Index

For an integer $i \geq 0$, define

$$[i] = \frac{q^i - 1}{q - 1} = q^{i-1} + q^{i-2} + \cdots + 1, \quad (2.10)$$

so that $[i]_{q=1} = i$. Recall that the *Stirling number* $S(r, \ell)$ is the number of set partitions of a set of size r into ℓ subsets. We now compute the sum that appears in Corollary 2.2.

Proposition 2.3. *For $r, n \in \mathbb{Z}_{>0}$,*

$$\sum_{a \in \{1, \dots, n\}^r} q^{\text{maj}(w_a)} = \sum_{\ell=1}^n S(r, \ell) [n] [n-1] \cdots [n-\ell+1].$$

Proof. We begin by classifying the permutations w_a that appear in the sum. For each sequence $a = (a_1, \dots, a_r) \in \{1, \dots, n\}^r$ we define $\text{shape}(a)$ to be the set partition of $\{1, \dots, r\}$ given by the rule,

$$i \sim j \text{ in } \text{shape}(a) \quad \text{if and only if} \quad a_i = a_j \text{ in } a.$$

We also let

$$D_{n,t} = \{ w = (w_1, \dots, w_n) \in S_n \mid w_1 < w_2 < \dots < w_t \},$$

be a distinct set of minimal-length coset representatives of S_n/S_t , where we naturally embed $S_t \subseteq S_n$ as the permutations of $\{1, \dots, t\}$. From this construction, we immediately have,

$$\text{if } \text{shape}(a) \text{ has } \ell \text{ parts} \quad \text{then} \quad w_a \in D_{n, n-\ell}.$$

For example, if $n = 6$, $r = 10$, and $a = (2, 1, 3, 1, 6, 2, 6, 1, 3, 1)$, then

$$\begin{aligned} a = (\underbrace{2, 1, 3, 1, 6, 2, 6, 1, 3, 1}_{\ell=4 \text{ distinct entries}}) &\Rightarrow w_a = (\underbrace{4, 5}_{n-\ell=2}, \underbrace{2, 6, 3, 1}_{\ell=4}) \in D_{6,2}, \\ \text{shape}(a) = \text{shape}(2, 1, 3, 1, 6, 2, 6, 1, 3, 1) &= \underbrace{\{1, 6\} \cup \{2, 4, 8, 10\} \cup \{3, 9\} \cup \{5, 7\}}_{\ell=4 \text{ parts}}. \end{aligned}$$

Note that the number of possible parts ℓ in $\text{shape}(a)$ is bounded both by the number r of subscripts and the number n of possible choices of a_i .

For a fixed set partition K with ℓ parts and a fixed permutation $w \in D_{n, n-\ell}$ we can easily reconstruct the unique sequence $a \in \{1, \dots, n\}^r$ with ℓ distinct entries such that $\text{shape}(a) = K$ and $w_a = w$. Thus, if we let $\mathcal{P}_r^\ell$ be the set partitions of $\{1, \dots, r\}$ with ℓ parts, then

$$\begin{aligned} \sum_{a \in \{1, \dots, n\}^r} q^{\text{maj}(w_a)} &= \sum_{\ell=1}^{\min(r, n)} \sum_{K \in \mathcal{P}_r^\ell} \sum_{\substack{a \in \{1, \dots, n\}^r \\ \text{shape}(a) = K}} q^{\text{maj}(w_a)} = \sum_{\ell=1}^{\min(n, r)} \sum_{K \in \mathcal{P}_r^\ell} \sum_{w \in D_{n, n-\ell}} q^{\text{maj}(w)} \\ &= \sum_{\ell=1}^{\min(n, r)} S(r, \ell) \sum_{w \in D_{n, n-\ell}} q^{\text{maj}(w)}, \end{aligned}$$

where the last equality comes from the fact that the Stirling number $S(r, \ell)$ is the number of partitions of $\{1, \dots, r\}$ into ℓ parts.

To finish the proof of the proposition, we will show that

$$\sum_{w \in D_{n,t}} q^{\text{maj}(w)} = [n][n-1] \cdots [t+1], \quad 0 \leq t < n. \quad (2.11)$$

The shape of a permutation w is the composition $\mu = (\mu_1, \dots, \mu_\ell)$ of n where μ_1 is the first position i where $w_i > w_{i+1}$, $\mu_1 + \mu_2$ is the next position i where $w_i > w_{i+1}$ and so on. The sum in (2.11) is over all partitions whose shape μ satisfies $\mu_1 \geq t$. An inversion in a permutation w is a pair (i, j) such that $i < j$ and $w_i > w_j$ and $\text{inv}(w)$ is the number of inversions in w . Foata and Schützenberger [FS] (see also [Lo, Theorem 11.4.4]) prove that the number of permutations of shape μ having m inversions equals the number of permutations of shape μ having m backsteps. Thus,

$$\sum_{w \in D_{n,t}} q^{\text{maj}(w)} = \sum_{w \in D_{n,t}} q^{\text{inv}(w)}. \quad (2.12)$$

Now, our coset representatives $D_{n,t}$ for S_n/S_t are chosen with minimal length, so if $u \in D_{n,t}$ and $v \in S_t$, then $\text{inv}(uv) = \text{inv}(u) + \text{inv}(v)$. Thus,

$$[n]! = \sum_{s \in S_n} q^{\text{inv}(s)} = \sum_{u \in D_{n,t}} \sum_{v \in S_t} q^{\text{inv}(uv)} = \sum_{u \in D_{n,t}} q^{\text{inv}(u)} \sum_{v \in S_t} q^{\text{inv}(v)} = \sum_{u \in D_{n,t}} q^{\text{inv}(u)} [t]!,$$

where the first and last equalities come from the well-known result of MacMahon (see [Sta1, Cor 1.3.10]) that $\sum_{w \in S_n} q^{\text{inv}(w)} = [n]!$. Equation (2.11) follows by dividing by $[t]!$ and replacing inv with maj . $\square$

For $n, r \in \mathbb{Z}_{>0}$, define,

$$d_{n,r}(q) = \sum_{\ell=1}^n S(r, \ell) [n][n-1] \cdots [n-\ell+1]. \quad (2.13)$$

The first few values of $d_{n,r}(q)$, for increasing r , are given by

$$\begin{aligned} d_{n,0}(q) &= 1, \\ d_{n,1}(q) &= [n], \\ d_{n,2}(q) &= [n]([n-1] + 1), \\ d_{n,3}(q) &= [n](1 + 3[n-1] + [n-1][n-2]), \\ d_{n,4}(q) &= [n](1 + 7[n-1] + 6[n-1][n-2] + [n-1][n-2][n-3]). \end{aligned}$$

When $q = 0$, we have $[j]_{q=0} = 1$, so $d_{n,r}(0) = \sum_{\ell=0}^n S(r, \ell)$, which equals the r th Bell number $B(r)$ if $n \geq r$ and which is the number of set partitions of $\{1, \dots, r\}$ into at most n subsets if $n < r$. When $q = 1$ the sum in Proposition 2.3 shows that $d_{n,r}(1)$ equals the cardinality of $\{1, \dots, n\}^r$, so $d_{n,r}(1) = n^r$. These polynomials are tantalizingly close to those in the following identity of Garsia and Remmel [GR, I.17]

$$\sum_{\ell=1}^r S(r, \ell; q) [n][n-1] \cdots [n-\ell+1] = [n]^r,$$

where $S(r, \ell; q)$ is a q -analog of the Stirling number $S(r, \ell)$. Like $d_{n,r}(q)$, these Garsia–Remmel polynomials specialize at $q = 1$ to n^r , but they are different at $q = 0$, since $[n]^r|_{q=0} = 1$.

The next Corollary follows immediately from Corollary 2.2 and Proposition 2.3.

Corollary 2.4. For $n, r \in \mathbb{Z}_{>0}$, we have

$$d_{n,r}(q) = \sum_{\ell=1}^n S(r, \ell) [n][n-1] \cdots [n-\ell+1] = \sum_{\lambda \in \Lambda_r^n} f_n^\lambda(q) m_r^\lambda.$$

3 A Basis for the $\mathcal{IR}$ Module for $\mathrm{GL}_n(\mathbb{F}_q)$


In this section we construct a module $\mathcal{IR}_q^r$ for the finite general linear group $\mathrm{GL}_n(\mathbb{F}_q)$ using r iterations of Harish-Chandra restriction and induction. We find a basis for $\mathcal{IR}_q^r$ that is indexed by q -set partitions of $\{1, \dots, r\}$. It is easy to see that the number of these is the polynomial $d_{n,r}(q)$ which appeared in Section 2, and so $\dim(\mathcal{IR}_q^r) = d_{n,r}(q)$. The module $\mathcal{IR}_q^r$ is the defining space for the q -partition algebra, which will be analyzed in a subsequent paper by T. Halverson, A. Ram, and N. Thiern.

3.1 A family of q -analogues to set partitions

Let

$$\mathbb{Z}_n^r = \{(k_1, k_2, \dots, k_r) \in \mathbb{Z}^r \mid 0 \leq k_1, k_2, \dots, k_r \leq n-1\},$$

which we can think of as a configuration of boxes stacked into an $(n-1) \times r$ rectangle. That is, $(k_1, k_2, \dots, k_r)$ denotes the collection of boxes with k_j boxes stacked in the j th column. For example,




Let

$$\mathcal{P}_{n \times r} = \{(k_1, k_2, \dots, k_r) \in \mathbb{Z}_n^r \mid k_1 = 0, k_j = h \text{ implies } k_i = h-1 \text{ for some } i < j\}.$$

We have a surjection

$$\begin{array}{ccc} \mathbb{Z}_n^r & \longrightarrow & \mathcal{P}_{n \times r} \\ k & \mapsto & k^*, \end{array} \quad \text{where } k_1^* = 0 \text{ and } k_j^* = \min \{k_j, \max\{k_1^* + 1, k_2^* + 1, \dots, k_{j-1}^* + 1\}\},$$

which sends




We will refer to k^* as the $*$ -height of k .

There is a bijection,

$$\begin{array}{ccc} \mathcal{P}_{n \times r} & \longleftrightarrow & \left\{ \begin{array}{l} \text{Set partitions of } \{1, 2, \dots, r\} \\ \text{with at most } n \text{ parts} \end{array} \right\} \\ k = (k_1, k_2, \dots, k_r) & \mapsto & K_k, \end{array}$$


where i and j are in the same part of K_k if and only if $k_i = k_j$. That is,



To obtain q -analogues, fill the boxes with elements of $\mathbb{F}_q$. Let

$$\mathbb{Z}_n^r(q) = \{((k_1, a^{(1)}), (k_2, a^{(2)}), \dots, (k_r, a^{(r)})) \mid (k_1, k_2, \dots, k_r) \in \mathbb{Z}_n^r, a^{(j)} \in \mathbb{F}_q^{k_j}\},$$


For example,



The q -analogue of set partitions of $\{1, 2, \dots, r\}$ with at most n parts is the set

$$\mathcal{P}_{n \times r}(q) = \{((k_1, a^{(1)}), (k_2, a^{(2)}), \dots, (k_r, a^{(r)})) \mid (k_1, k_2, \dots, k_r) \in \mathbb{Z}_n^r, a^{(j)} \in \mathbb{F}_q^{k_j - k_j^*}\}.$$

For example,



where the boxes labeled by $*$ give the $*$ -height for the associated element in $\mathbb{Z}_n^r$.

An n -restricted q -set partition of $\{1, 2, \dots, r\}$ is an element of $\mathcal{P}_{n \times r}(q)$. Given a set partition K_h with ℓ parts, there are

$$[n][n-1] \cdots [n-\ell+1]$$

different n -restricted q -set partitions of $\{1, 2, \dots, r\}$ with $*$ -height h . Thus,

$$|\mathcal{P}_{n \times r}(q)| = \sum_{\ell=1}^n S(r, \ell) [n][n-1] \cdots [n-\ell+1] = d_{n,r}(q),$$

where $d_{n,r}(q)$ is defined in (2.13). By the constructions of this section, we also easily obtain the specializations,

$$\begin{aligned} |\mathcal{P}_{n \times r}(0)| &= B(r), & \text{for } n \geq r, \\ |\mathcal{P}_{n \times r}(1)| &= |\mathbb{Z}_n^r(1)| = n^r, \end{aligned}$$

where $B(r)$ is the r th Bell number.

3.2 The Chevalley group $GL_n(\mathbb{F}_q)$

The general linear group $G_n = GL_n(\mathbb{F}_q)$ has a double coset decomposition given by

$$G_n = \bigsqcup_{w \in S_n} U_B w B_n, \quad (3.1)$$

where S_n is the subgroup of permutation matrices, and

$$B_n = \left\{ \begin{pmatrix} * & & * \\ & \ddots & \\ 0 & & * \end{pmatrix} \right\} \subseteq G_n \quad \text{and} \quad U_B = \left\{ \begin{pmatrix} 1 & & * \\ & \ddots & \\ 0 & & 1 \end{pmatrix} \right\} \subseteq B_n$$

are the subgroups of upper-triangular matrices and unipotent upper-triangular matrices, respectively. For $1 \leq i < j \leq n$ and $a \in \mathbb{F}_q$, let $x_{ij}(a) \in U_B$ be the matrix with a in the (i, j) th position, ones on the diagonal, and zeroes everywhere else. Note that for $i < j$, $k < l$, $a, b \in \mathbb{F}_q$,

$$x_{ij}(a)x_{kl}(b) = \begin{cases} x_{kl}(b)x_{il}(ab)x_{ij}(a), & \text{if } j = k, \\ x_{kl}(b)x_{kj}(-ab)x_{ij}(a), & \text{if } i = l, \\ x_{ij}(a+b), & \text{if } i = k, j = l, \\ x_{kl}(b)x_{ij}(a), & \text{otherwise.} \end{cases} \quad (3.2)$$

For $w \in S_n$, we have

$$x_{ij}(a)w = wx_{w^{-1}(i)w^{-1}(j)}(a). \quad (3.3)$$

Let

$$L_n = \left\{ \left(\begin{array}{c|c} G_1 & 0 \\ \hline 0 & G_{n-1} \end{array} \right) \right\} \subseteq G_n, \quad U_n = \left\{ \left(\begin{array}{c|ccc} 1 & * & \cdots & * \\ \hline 0 & & & \text{Id}_{n-1} \end{array} \right) \right\} \subseteq G_n,$$

and

$$P_n = L_n U_n = \left\{ \left(\begin{array}{c|ccc} G_1 & * & \cdots & * \\ \hline 0 & & & G_{n-1} \end{array} \right) \right\}.$$

For $1 \leq k \leq n-1$, let

$$w_k = s_k s_{k-1} \cdots s_1,$$

where s_i is the simple reflection that switches i and $i+1$. By convention, $w_0 = 1$. Note that the $w_k, 0 \leq k \leq n-1$, give a set of minimal-length coset representatives for $S_n/(S_1 \times S_{n-1})$. For $1 \leq k \leq n-1$ and $a = (a_1, a_2, \dots, a_k) \in \mathbb{F}_q^k$, let

$$w_k(a) = s_k(a_k)s_{k-1}(a_{k-1}) \cdots s_1(a_1), \quad \text{where} \quad s_i(a_i) = x_{i,i+1}(a_i)s_i.$$

Then the decomposition

$$G_n = \bigsqcup_{\substack{0 \leq k \leq n-1 \\ a \in \mathbb{F}_q^k}} w_k(a)P_n \quad (3.4)$$

follows from (3.1) and (3.3).

3.3 Harish-Chandra Restriction and Induction

To make the notation more manageable, in this section we will assume that n is fixed and drop the subscripts in G_n, P_n, U_n, L_n . Let

$$e_U = \frac{1}{|U|} \sum_{u \in U} u,$$

so that $xe_U = e_U = e_Ux$ for all $x \in U$. Since U is a normal subgroup in P , there is a surjection $P \rightarrow P/U \cong L$, which gives rise to adjoint functors, called inflation and deflation, respectively,

$$\begin{aligned} \text{Inf}_L^P : \quad & \begin{array}{ccc} \{\text{Left } L\text{-modules}\} & \longrightarrow & \{\text{Left } P\text{-modules}\}, \\ V & \mapsto & e_U V \end{array} \\ \text{Def}_L^P : \quad & \begin{array}{ccc} \{\text{Left } P\text{-modules}\} & \longrightarrow & \{\text{Left } L\text{-modules}\}. \\ V & \mapsto & e_U V \end{array} \end{aligned}$$

By composing with induction and restriction, we obtain two functors

$$\begin{aligned} \text{Indf}_L^P : \quad & \begin{array}{ccccc} \{\text{Left } L\text{-modules}\} & \longrightarrow & \{\text{Left } P\text{-modules}\} & \longrightarrow & \{\text{Left } G\text{-modules}\}, \\ V & \mapsto & e_U V & \mapsto & \mathbb{C}G \otimes_{\mathbb{C}P} e_U V \end{array} \\ \text{Resf}_L^P : \quad & \begin{array}{ccccc} \{\text{Left } G\text{-modules}\} & \longrightarrow & \{\text{Left } P\text{-modules}\} & \longrightarrow & \{\text{Left } L\text{-modules}\}. \\ V & \mapsto & V & \mapsto & e_U V \end{array} \end{aligned}$$

Let $\mathbb{1}$ denote the trivial module of G . Define the G -module

$$\mathcal{IR}_q^r = (\text{Indf}_L^G \text{Resf}_L^G)^r(\mathbb{1}), \quad \text{for } r \geq 0, \quad (3.5)$$

and the L -module

$$\mathcal{IR}_q^{r+\frac{1}{2}} = \text{Resf}_L^G (\text{Indf}_L^G \text{Resf}_L^G)^r(\mathbb{1}), \quad \text{for } r \geq 0. \quad (3.6)$$

3.4 A Basis for $\mathcal{IR}_q^r$

Let

$$\otimes_U = \otimes_{\mathbb{C}P} e_U$$

denote tensoring over $\mathbb{C}P$ and multiplying by e_U . By construction it is clear that

$$\begin{aligned} \mathcal{IR}_q^r &= \mathbb{C}\text{-span}\{g_1 \otimes_U g_2 \otimes_U \cdots \otimes_U g_r \otimes \mathbb{1} \mid g_1, g_2, \dots, g_r \in G\} \\ &= \mathbb{C}\text{-span}\{w_{k_1}(a^{(1)}) \otimes_U \cdots \otimes_U w_{k_r}(a^{(r)}) \otimes \mathbb{1} \mid 0 \leq k_1, \dots, k_r \leq n-1, a^{(m)} \in \mathbb{F}_q^{k_m}\}, \\ \mathcal{IR}_q^{r+1/2} &= \mathbb{C}\text{-span}\{e_U g_1 \otimes_U g_2 \otimes_U \cdots \otimes_U g_r \otimes \mathbb{1} \mid g_1, g_2, \dots, g_r \in G\} \\ &= \mathbb{C}\text{-span}\{e_U w_{k_1}(a^{(1)}) \otimes_U \cdots \otimes_U w_{k_r}(a^{(r)}) \otimes \mathbb{1} \mid 0 \leq k_1, \dots, k_r \leq n-1, a^{(m)} \in \mathbb{F}_q^{k_m}\}. \end{aligned}$$

However, these sets are generally not linearly independent. The following lemma characterizes when two vectors are equal.

Lemma 3.1. *Fix $0 \leq k_1, k_2, \dots, k_l \leq n-1$ and $a^{(m)} \in \mathbb{F}_q^{k_m}$. Let $1 \leq i < j < n$ and $t \in \mathbb{F}_q^\times$. Then*

$$w_{k_1}(a^{(1)}) \otimes_U \cdots \otimes_U w_{k_l}(a^{(l)}) \otimes_U x_{ij}(t) = w_{k_1}(a^{(1)}) \otimes_U \cdots \otimes_U w_{k_l}(a^{(l)}) \otimes_U 1 \quad (3.7)$$

if and only if $i = 1$ or there exists $1 < m \leq l$ such that

$$w_{k_m} w_{k_{m+1}} \cdots w_{k_l} \quad \text{sends } i \text{ to } 1.$$

Proof. Note that for $i = 1$, $e_U x_{1j}(t) = e_U$ and for $i > 1$, $e_U x_{ij}(t) = x_{ij}(t)e_U$ (since U is normal in U_B). It therefore follows from (3.2) and (3.3) that for $0 \leq k \leq n-1$ and $a \in \mathbb{F}_q^k$,

$$w_k(a) \otimes_U x_{ij}(t) = \begin{cases} x_{w_k(i)j}(t)w_k(a) \otimes_U 1, & \text{if } k+1 < i < j, \\ x_{w_k(i),j}(t)w_k(a) \otimes_U 1, & \text{if } 1 < i \leq k+1 < j, \\ x_{w_k(i),j-1}(t)x_{w_k(i),k+1}(-a_{j-1}t)w_k(a) \otimes_U 1, & \text{if } 1 < i < j \leq k+1, \\ w_k(a) \otimes_U 1, & \text{if } i = 1. \end{cases} \quad (3.8)$$

If $i = 1$, then (3.7) follows. If $i > 1$ and there exists $1 < m < l$ such that $w_{k_m} w_{k_{m+1}} \cdots w_{k_l}$ sends i to 1, then using (3.8) we can push $x_{ij}(t)$ left past the $w_k(a)$ so that

$$e_U w_{k_m}(a^{(m)}) \otimes_U \cdots \otimes_U w_{k_l}(a^{(l)}) \otimes_U x_{ij}(t) = e_U u w_{k_m}(a^{(m)}) \otimes_U \cdots \otimes_U w_{k_l}(a^{(l)}) \otimes_U 1,$$

where $u \in U$ (since $w_{k_m} \cdots w_{k_l}(i) = 1$). But $e_U u = e_U$, giving (3.7).

Conversely, note that $w_k(a) \otimes_U x_{ij}(t) = w_k(a) \otimes_U 1$ if and only if $x_{ij}(t) \in U$. Thus, if (3.7) is true, then as we push $x_{ij}(t)$ to the left, at some point we must have

$$\begin{aligned} & w_{k_1}(a^{(1)}) \otimes_U \cdots \otimes_U w_{k_l}(a^{(l)}) \otimes_U x_{ij}(t) \\ &= w_{k_1}(a^{(1)}) \otimes_U \cdots \otimes_U w_{k_{m-1}}(a^{(m-1)}) \otimes_U u w_{k_m}(a^{(m)}) \otimes_U \cdots \otimes_U w_{k_l}(a^{(l)}) \otimes_U 1. \end{aligned}$$

for some $u \in U$. By (3.8) this can only happen if $w_{k_m} w_{k_{m+1}} \cdots w_{k_l}(i) = 1$ for some $1 < m \leq l$. $\square$

Combinatorially, we associate a column of labeled boxes to $w_k(a)$.

$$\begin{array}{|c|} \hline \\ \hline a_k \\ \hline \vdots \\ \hline a_2 \\ \hline a_1 \\ \hline \end{array} \longleftrightarrow w_k(a) = s_k(a_k) \cdots s_2(a_2) s_1(a_1). \quad (3.9)$$

We obtain vectors in $\mathcal{IR}_q^r$ by labeling r stacks of boxes. For example,

$$\longleftrightarrow w_1(a) \otimes_U w_4(b) \otimes_U w_0 \otimes_U w_2(c) \otimes_U w_2(d) \otimes_U w_6(e) \otimes_U w_3(f) \otimes \mathbb{1}.$$

Lemma 3.1 implies that not all choices of the vectors a, b, c, d, e, f will give different basis vectors of W_q^7 . In our example, any change to the *-ed values in

will not change the vector in $\mathcal{IR}^i$. That is, given an element of $\mathbb{Z}_n^r(q)$, Lemma 3.1 implies that the $*$ -height determines which entries can have arbitrary values (see Section 3.1). In particular, if the k_l th entry

$$\begin{aligned} & w_{k_1}(a^{(1)}) \otimes_U \cdots \otimes_U w_{k_{l-1}}(a^{(l-1)}) \otimes_U w_{k_l}(b) \otimes_U \cdots \\ & = w_{k_1}(a^{(1)}) \otimes_U \cdots \otimes_U w_{k_{l-1}}(a^{(l-1)}) \otimes_U x_{1,k_l+1}(b_1) \cdots x_{k_l,k_l+1}(b_{k_l}) w_{k_l} \otimes_U \cdots \end{aligned}$$

has $*$ -height h then for each $1 \leq i \leq h$, there exists $1 < m_i \leq l - 1$ such that

$$w_{k_{m_i}} \cdots w_{k_{l-1}}(i) = 1.$$

By Lemma 3.1 these entries can be replaced by arbitrary entries, and for these entries we average over all possible choices. Thus, for each element in $\mathcal{P}_{n \times r}(q)$, we obtain a basis vector. Specifically, for $0 \leq k^* \leq k \leq n-1$ and $a = (a_1, \dots, a_{k-k^*}) \in \mathbb{F}_q^{k-k^*}$, associate

$$\begin{array}{|c|} \hline \\ \hline \\ \hline \\ \vdots \\ \hline a_1 \\ \hline * \\ \hline \vdots \\ \hline * \\ \hline \end{array} \longleftrightarrow w_k^*(a) = s_k(a_{k-k^*}) \cdots s_{k^*+1}(a_1) \frac{1}{q^{k^*}} \sum_{b \in \mathbb{F}_q^{k^*}} s_{k^*}(b_{k^*}) \cdots s_1(b_1). \quad (3.10)$$

For $K = ((k_1, a^{(1)}), \dots, (k_r, a^{(r)})) \in \mathcal{P}_{n \times r}(q)$, let

$$v_K = \overset{*}{w}_{k_1}(a^{(1)}) \otimes_U \overset{*}{w}_{k_2}(a^{(2)}) \otimes_U \cdots \otimes_U \overset{*}{w}_{k_r}(a^{(r)}) \in \mathbb{C}G \otimes_U \mathbb{C}G \otimes_U \cdots \otimes_U \mathbb{C}G \otimes_U \mathbf{1}_n.$$

For example,

$$\begin{aligned}
v \begin{array}{|c|c|c|c|} \hline & & g & \\ \hline & d & f & \\ \hline & c & e & \\ \hline & b & * & * \\ \hline a & * & * & * \\ \hline \end{array} &= x_{12}(a)w_1 \otimes x_{45}(d)x_{35}(c)x_{25}(b) \frac{1}{q} \sum_{t \in \mathbb{F}_q} x_{15}(t)w_4 \otimes 1 \otimes \frac{1}{q^2} \sum_{s,t \in \mathbb{F}_q} x_{23}(s)x_{13}(t)w_2 \\
&\otimes \frac{1}{q^2} \sum_{s,t \in \mathbb{F}_q^\times} x_{23}(s)x_{13}(t)w_2 \otimes x_{67}(g)x_{57}(f)x_{47}(e) \frac{1}{q^3} \sum_{r,s,t \in \mathbb{F}_q} x_{37}(r)x_{27}(s)x_{17}(t)w_6 \\
&\otimes \frac{1}{q^3} \sum_{r,s,t \in \mathbb{F}_q} x_{34}(r)x_{24}(s)x_{14}(t)w_3 \otimes \mathbb{1}_n.
\end{aligned}$$

Lemma 3.1 and the following discussion imply that the v_K are linearly independent, so we have proved the first part of the following theorem.

Theorem 3.2. *Let $r \in \mathbb{Z}_{\geq 0}$. Then*

(a) The G -module $\mathcal{IR}_q^r$ has a basis given by

$$\{v_K \mid K \in \mathcal{P}_{n \times r}(q)\},$$

and thus $\dim(\mathcal{IR}_q^r) = d_{n,r}(q)$.

(b) The L -module $W_q^{r+1/2}$ has a basis given by

$$\{v_K \mid K \in \mathcal{P}_{n \times r+1}(q) \text{ with } k_1 = 0\},$$

and thus $\dim(\mathcal{IR}_q^r) = d_{n,r+1}(q)/[n]$.


To prove Theorem 3.2 (b), it suffices to characterize what happens in $e_U \mathcal{IR}_q^r$. Let

$$\begin{array}{ccc} \mathcal{P}_{n \times r}(q) & \longrightarrow & \{K \in \mathcal{P}_{n \times r+1}(q) \mid k_1 = 0\} \\ K & \mapsto & \tilde{K} \end{array}$$

be the surjective function given by the following algorithm.

- (1) Add an empty column to the left side of K and set $m = 1$,
- (2) If the resulting diagram is in $\mathcal{P}_{n \times r+1}(q)$, stop. Else set $m := m + 1$.
- (3) If column m has an unstarred box, then replace the bottom unstarred entry by $*$. Go to step (2).

For example, we get



Lemma 3.3. Let $K \in \mathcal{P}_{n \times k}(q)$. Then $e_U v_K = v_{\tilde{K}}$.

Proof. Apply Lemma 3.1 to the vector

$$w_0 \otimes_U v_K$$

to obtain the statement of this lemma. $\square$

Proof of Theorem 3.2 (b). We have that

$$\begin{aligned} \mathcal{IR}_q^{r+1/2} &= e_U \mathcal{IR}_q^r \\ &= \mathbb{C}\text{-span}\{e_U v_K \mid K \in \mathcal{P}_{n \times r}(q)\} \\ &= \mathbb{C}\text{-span}\{v_{\tilde{K}} \mid \mathcal{P}_{n \times r}(q)\}, \end{aligned}$$

and by Lemma 3.1, the vectors in the last set are linearly independent. $\square$

4 Group action on $\mathcal{IR}^r$

In general,

$$gw_k(a) \otimes_U v = w_l(b) \otimes_U pv, \quad \text{where } gw_k(a) = w_l(b)p.$$

Thus, globally the matrix of g is the matrix of g acting by left multiplication on G/P . The group G has generators given by

$$\{x_{ij}(t) \mid 1 \leq i < j \leq n, t \in \mathbb{F}_q\} \cup \{s_1, s_2, \dots, s_{n-1}\} \cup \{h_k(t) \mid 1 \leq k \leq n, t \in \mathbb{F}_q^\times\},$$

where $h_k(t)$ is the identity matrix with the k th diagonal 1 replaced by t . The generators of G act on $\mathcal{IR}_q^r$ in the following way:

$$\begin{aligned} s_i w_k(a) \otimes_U v &= \begin{cases} w_k(a) \otimes_U s_i v, & \text{if } i > k+1, \\ w_{k+1}(a_1, a_2, \dots, a_k, 0) \otimes_U v, & \text{if } i = k+1, \\ w_{k-1}(a_1, \dots, a_{k-1}) \otimes_U v, & \text{if } i = k, a_k = 0, \\ w_k(a_1, \dots, a_{k-1}, a_k^{-1}) \otimes_U h_{k+1}(-a_k^{-1})v, & \text{if } i = k, a_k \neq 0, \\ w_k(a_1, \dots, a_{i-1}, a_{i+1}, a_i, a_{i+2}, \dots, a_k) \otimes_U s_{i+1}v, & \text{if } i < k. \end{cases} \\ h_j(b) w_k(a) \otimes_U v &= \begin{cases} w_k(a) \otimes_U h_j(b)v, & \text{if } j > k+1, \\ w_k(a_1 b^{-1}, \dots, a_k b^{-1}) \otimes_U v, & \text{if } j = k+1, \\ w_k(a_1, \dots, a_{j-1}, b a_j, a_{j+1}, \dots, a_k) \otimes_U h_{j+1}(b)v, & \text{if } i < k+1. \end{cases} \\ x_{ij}(b) w_k(a) \otimes_U v &= \begin{cases} w_k(a) \otimes_U x_{ij}(b)v, & \text{if } i > k+1, \\ w_k(a) \otimes_U x_{k+1,j}(-a_k b) \cdots x_{2,j}(-a_1 b)v, & \text{if } i = k+1 \neq 1, \\ w_k(a) \otimes_U v, & \text{if } i = k+1 = 1, \\ w_k(a) \otimes_U x_{i+1,j}(b)v, & \text{if } i < k+1 < j, \\ w_k(a_1, \dots, a_{i-1}, a_i + b, a_{i+1}, \dots, a_k) \otimes_U v, & \text{if } i < k+1 = j, \\ w_k(a_1, \dots, a_{i-1}, a_i + b a_j, a_{i+1}, \dots, a_k) \otimes_U x_{i+1,j+1}(b)v, & \text{if } j < k+1. \end{cases} \end{aligned}$$

References

- [BW] J. Birman and H. Wenzl, Braids, link polynomials, and a new algebra, *Trans. Amer. Math. Soc.*, **313** (1989), 249-273.
- [CDDSY] W. Chen, E. Deng, R. Du, R. Stanley, and C. Yan, Crossings and nestings of matchings and partitions, *Trans. Amer. Math. Soc.* **359** (2007), 1555–1575.

- [FS] D. Foata and M-P. Schützenberger, Major index and inversion number of permutations, *Math. Nachr.*, **83** (1978), 143–159.
- [GR] A. Garsia and J. Remmel, Q -counting rook configurations and a formula of Frobenius, *J. Combin. Theory Ser. A*, **41** (1986), 246–275.
- [HL] T. Halverson and T. Lewandowski, RSK insertion for set partitions and diagram algebras *Electron. J. Combin.*, **11** (2004/06), 24.
- [HR] T. Halverson and A. Ram, Partition Algebras, *European J. Combinatorics*, **26** (2005), 869–921.
- [Ji] M. Jimbo, A q -analog of $U(\mathfrak{gl}(N + 1))$, Hecke algebra, and the Yang-Baxter equation, *Lett. Math. Phys.*, **11** (1986), 247–252.
- [Jo] V. F. R. Jones, The Potts model and the symmetric group, in: *Subfactors: Proceedings of the Taniguchi Symposium on Operator Algebras (Kyuzeso, 1993)*, World Sci. Publishing, River Edge, NJ, 1994, 259–267.
- [Lo] M. Lothaire, *Algebraic combinatorics on words*, Encyclopedia of Mathematics and its Applications, **90** Cambridge University Press, Cambridge, 2002.
- [Mac] I. Macdonald, *Symmetric Functions and Hall Polynomials*, 2nd ed., Oxford Univ. Press, New York, 1995.
- [Mar1] P. Martin, Temperley-Lieb algebras for nonplanar statistical mechanics—the partition algebra construction, *J. Knot Theory Ramifications* **3** (1994), 51–82.
- [Mar2] P. Martin, The structure of the partition algebras, *J. Algebra* **183** (1996), 319–358.
- [Mur] J. Murakami, The Kauffman polynomial of links and representation theory, *Osaka J. Math.*, **24** (1987), 745–758
- [Sta1] R. Stanley, *Enumerative Combinatorics, Vol. 1*, Cambridge University Press, Cambridge, 1997.
- [Sta2] R. Stanley, *Enumerative Combinatorics, Vol. 2*, Cambridge University Press, Cambridge, 1999.